

LiBeB: High and low energy cosmic ray production and comparison with ν induced nucleosynthesis in SNII

Reuven Ramaty

Goddard Space Flight Center, Greenbelt, MD 20771, USA

Hubert Reeves

Institut d'Astrophysique de Paris, Paris F-75014, France

Richard E. Lingenfelter

University of California San Diego, LaJolla, CA 92093, USA

and

Benzion Kozlovsky

Sackler Faculty of Exact Sciences, Tel Aviv University, Israel

Abstract

We present new calculations of LiBeB production by accelerated particles with various compositions and energy spectra ranging from low energies to relativistic energies, and various ambient medium metallicities ($Z/Z_{\odot}$). The observed, essentially constant, Be/Fe ratio as a function of $Z/Z_{\odot}$ can be best understood if the metallicity of the accelerated particles (high energy or low energy) does not vary much with $Z/Z_{\odot}$. This could be achieved if the particles are accelerated directly from the ejecta of Type II supernovae (SNII) and not from the interstellar medium. Using the observed Be/Fe and the fact that most of the Fe at low $Z/Z_{\odot}$ is from SNII, we derive the energy content in accelerated particles per SNII (a few times 10^{50} erg). We show that additional ^{11}B production by neutrinos is consistent with the available data, allowing a neutrino yield from SNII less than or equal to the nominal published yields. We further show that the observed ^6Li at low $Z/Z_{\odot}$ suggests that the accelerated particles responsible for the LiBeB at such $Z/Z_{\odot}$ are confined to low energies and have a relatively high He/O abundance.

In press: Nuclei in the Cosmos 1996, Nuclear Physics, Conference Proceedings

1 INTRODUCTION

Observations of stars of various metallicities ($Z/Z_{\odot}=(\text{Fe}/\text{H})/(\text{Fe}/\text{H})_{\odot}$) have shown that the abundance ratios B/Fe and Be/Fe [1] do not vary appreciably over a broad range of $Z/Z_{\odot}$ (see [2]). This near constancy implies that the ratios of the B and Be production rates to the Fe production rate must remain essentially constant as a function of metallicity. Moreover, as long as this constancy is maintained and destruction effects (due to astration) do not play a major role, the production and inventory ratios are equal. For $Z/Z_{\odot}$ less than about 0.1, most of the Fe is thought to be produced in Type II supernovae (SNII), with Type I supernovae not yet contributing significantly [3]. Limiting our considerations to this range of metallicities, the constancy of B/Fe and Be/Fe strongly suggest that B and Be are also produced by SNIIs. The accepted paradigm for the origin of most of the Be is production in spallation reactions of accelerated particles (p , α , CNO) deriving their energy from supernovae. B, in addition to such reactions, is also produced in neutrino interactions in SNIIs [4, 5, 6]. The observed Be abundance as a function of metallicity can be approximated by $\text{Be}/\text{H} \simeq 5.2 \times 10^{-11} (\text{Fe}/\text{H})/(\text{Fe}/\text{H})_{\odot}$ [2], implying that $\text{Be}/\text{Fe} \simeq 1.7 \times 10^{-6}$. The relatively well known Fe yield per SNII ($\sim 0.1 M_{\odot}$ based on SN87A observations) then provides a direct determination of the accelerated particle-produced Be per SNII ($\sim 3.6 \times 10^{48}$ atoms), independent of the details of the spallation reactions. The total energy per SNII imparted to the accelerated particles to produce the required amount of Be, as well as the accompanying production of the other light isotopes, do depend on the accelerated particle composition and spectrum, the ambient medium composition and the path length X_{esc} available to the accelerated particles for producing nuclear reactions.

Previous calculation of LiBeB production by accelerated particles [7, 8] incorporated various compositions and energy spectra but used different codes for spectra extending to relativistic energies (hereafter HECR) or spectra confined to low energies (hereafter LECR). Furthermore, the HECR calculations at epochs of low metallicity implicitly assumed that the composition of the HECR is the same as that of the ambient medium [9, 10, 7]. In addition, secondary reaction chains (e.g. fast ^{12}C on ambient H producing ^{11}B and the subsequent interaction of the fast ^{11}B to produce ^{10}B or lighter isotopes), which are quite important when HECR with solar or higher metallicity are interacting with a low metallicity ambient medium, have been ignored. We

have developed a new code that employs updated cross sections [11, 12, 13] and incorporates all the possible secondary chains for source particles up to ^{16}O . We calculate LiBeB production for a variety of accelerated particle spectra (LECR to HECR), ambient medium metallicities $10^{-3} < Z/Z_{\odot} < 1$, and various accelerated particle compositions. By normalizing the LiBeB yields to the energy content of the source accelerated particles, we can directly determine the SNII particle acceleration efficiency required to produce the observed Be/Fe. We next combine the accelerated particle-produced and ν -produced ^{11}B and set constraints on the contribution of the ν process on B production. Finally, we consider the implications of the ^6Li observations at low metallicity [17].

2 ANALYSIS

We assume that the ambient composition is solar scaled by a variable metallicity. For the fast particle composition we use the following: (i) ejecta of Type II supernovae (12-40 $M_{\odot}$ progenitors) for various metallicities [5] averaged over an IMF proportional to $M^{-2.7}$ (hereafter SNII); (ii) current epoch cosmic ray source abundances [14, 15] (hereafter CRS); (iii) ambient medium abundances, i.e. solar system abundances scaled with metallicity (hereafter SSz); and (iv) the ejecta of a supernova from a 60 $M_{\odot}$ progenitor [16] (hereafter SN60). The SNII abundances of the most relevant isotopes, ^1H , ^4He , ^{12}C and ^{16}O , are weakly dependent on $Z/Z_{\odot}$ and are approximately proportional to 100:20:0.2:1, respectively. The CRS and SN60 abundances are taken independent of $Z/Z_{\odot}$ and the above proportions are 175:25:0.8:1 and 0:0.7:0.5:1, respectively. Thus, CRS is not too different from SNII except that it is more abundant in ^{12}C ; for SN60, H and He are absent or very under abundant, and, as for CRS, ^{12}C is more abundant than for SNII. We employ a shock acceleration spectrum with a high energy cutoff, $(p^{-s}/\beta)e^{-E/E_0}$, except for SN60 for which, as before [8], we assume that above 50 MeV/nuc the spectrum decreases as E^{-10} ; p and E are momentum and kinetic energy, both per nucleon, and $c\beta$ is particle velocity. The SN60 model, because of its relatively high C/O and sharp high energy cutoff, produces the highest B/Be and $^{11}\text{B}/^{10}\text{B}$ without violating energy constraints (see below). We assume that X_{esc} is energy independent, and present results for $X_{esc}=10 \text{ g cm}^{-2}$, a leaky box model, and $X_{esc} \rightarrow \infty$, a closed galaxy model. The neglect of the

energy dependence affects our results only marginally. The incorporation of the secondary reaction chains can lead to contributions as high as 60% of the total production in some cases.

The nuclear reactions are accompanied by energy loss due to Coulomb interactions of the accelerated particles with the ambient medium (assumed to be neutral). Fig. 1 shows the Be production per deposited energy. The solid curves are for SNII, the dashed line is for SSz and the dashed line with diamonds is for SN60. The values for CRS are practically identical to those for SNII with $E_0=10\text{GeV/n}$. All curves with given values of E_0 are for the exponential cutoff at the indicated E_0 . We see that for SNII and SN60, for which the metallicities of the accelerated particles are nearly independent of $Z/Z_\odot$, $Q(\text{Be})/W$ is practically independent of the ambient medium metallicity; this, however, is not the case for SSz for which the metallicity of the fast particles increases with increasing $Z/Z_\odot$. $Q(\text{Be})/W$ decreases rapidly for $E_0 < 50\text{MeV/n}$; we show values as a function of $Z/Z_\odot$ for $E_0=10\text{MeV/n}$. Using the values plotted in Fig. 1 we can express the Be yield per SNII as $Q(\text{Be}) = 10^{51}W_{51}[Q(\text{Be})/W]$, where $10^{51}W_{51}\text{ erg}$ is the energy content in accelerated particles per SNII. As the total SNII mechanical energy is $\sim 10^{51}\text{ erg}$, W_{51} is essentially the acceleration efficiency. We assume that W_{51} is independent of $Z/Z_\odot$.

The calculated [5] ^{56}Fe yields, averaged over the IMF given above, range from about 0.09 to 0.14 $M_\odot$ per SNII. As already indicated, we adopt the constant value of 0.1, leading to a required Be production per SNII of 3.6×10^{48} atoms independent of $Z/Z_\odot$. Then for the SNII, CRS and SN60 compositions, for which $Q(\text{Be})/W$ is independent of $Z/Z_\odot$, $W_{51}[Q(\text{Be})/W] \simeq 3.6 \times 10^{-3}$. Requiring an acceleration efficiency less than about 0.3, $Q(\text{Be})/W \geq 1.2 \times 10^{-2}$. This effectively rules out cutoff energies $E_0 < 50\text{ MeV/n}$, as well as the SSz case. The latter is also ruled because of its dependence on $Z/Z_\odot$.

Fig. 2 shows $Q(\text{B})/Q(\text{Be})$, the accelerated particle produced B-Be ratio, for all cases except SSz. $Q(\text{B})/Q(\text{Be})$ is practically independent of $Z/Z_\odot$, implying that $(\text{B/Be})_{cr}$ can remain constant as a function of $Z/Z_\odot$. The calculated $Q(\text{B})/Q(\text{Be})$ ranges from about 10 to 30, depending on the accelerated particle spectrum and composition. ^{11}B can also be produced by ν spallation in supernovae [4, 5]. By averaging the calculated ^{11}B yields over the same IMF, we obtain a ν -induced ^{11}B production of $\simeq 6 \times 10^{-7} M_\odot$ per SNII, practically independent of metallicity. For the total B-to-Be inventory ratio we thus have

$$B/Be \simeq (B/Be)_{cr} + 3.0M_7, \quad (1)$$

where M_7 is the ν -produced ^{11}B in units of $10^{-7}M_\odot$. The second term in eq. (1) is simply determined by the numbers of ^{56}Fe and ν -produced ^{11}B per SNII, and the observed Be/Fe ratio. The observed B/Be is quite uncertain, mainly because of NLTE effects. Adopting the range $9 < B/Be < 34$ [18] for low metallicity stars, the maximum contribution of ν spallation, $M_7 \simeq 8$, is obtained for the minimal $(B/Be)_{cr} \simeq 10$ corresponding to production by accelerated particles extending to GeV energies in the closed galaxy model. This upper limit slightly exceeds the predicted ^{11}B production by neutrinos [5]. For the other accelerated particle compositions, $Q(B)/Q(Be)$ is larger and consequently the maximal M_7 is smaller. For example, for SNII with $E_0 = 50 \text{ MeV/n}$, $Q(B)/Q(Be) \simeq 18$ and hence $M_7 \leq 5$, a limit which is slightly lower than the predicted value.

Fig. 3 shows $Q(^{11}B)/Q(^{10}B)$ together with the meteoritic B isotopic ratio of about 4. This isotopic ratio cannot be reproduced by HECR which yield an isotopic ratio less than about 2.5 (Fig. 3). The addition of the ν -produced ^{11}B can provide the extra ^{11}B . Using eq. (1) we obtain $^{11}B/^{10}B = [3M_7/(B/Be)_{cr}] [(^{11}B/^{10}B)_{cr} + 1] + (^{11}B/^{10}B)_{cr}$, where $^{11}B/^{10}B$ includes both the accelerated particle and ν -produced ^{11}B . If for the low metallicity stars $^{11}B/^{10}B$ equals the solar system value of 4, then for the various models $0 < M_7 < 2$, i.e. ranging from a negligible ν contribution to less than about 30% of the predicted value.

Fig. 4 shows $Q(^6Li)/Q(Be)$. For the compositions that we use, this ratio also remains constant as a function of $Z/Z_\odot$ allowing us to compare production and inventory abundance ratios. A value of 0.05 ± 0.02 for $^6Li/Li$ was observed at $Z/Z_\odot \simeq 10^{-2.4}$ [17]. The error bar (SLN) shown at this metallicity is based on this observation, and on the observed Li/H (Spite plateau, error $\pm 47\%$) and Be/H at $Z/Z_\odot \simeq 10^{-2.4}$, for which we estimate an error of $\pm 50\%$. We see that SNII with $E_0 \simeq 50 \text{ MeV/n}$ produces more ^6Li relative to Be than both HECR and SN60. HECR underproduce ^6Li because the $\alpha - \alpha$ cross section is relatively low at high energies; and SN60 underproduces ^6Li because of the low α particle abundance. Although the large error bar of the SLN data does not allow us to discriminate between the models, these data favor SNII with $E_0 \simeq 50 \text{ MeV/n}$ rather than either HECR or SN60. On the other hand, the meteoritic $^6Li/Be$ at $Z/Z_\odot = 1$ (Fig. 4) is consistent with these models suggesting that the $LiBeB$ at the time of the formation of the

solar system resulted from more than one accelerated particle component. We mention in addition that the high $Q(^6\text{Li})/Q(\text{Be})$ implied by SNII with $E_0 \simeq 50 \text{ MeV/n}$ tends to overproduce Li/H at $Z/Z_\odot$ greater than about 0.1. To overcome this problem, the SN60 model was used in a recent galactic abundance evolution study [6]. But as we see from Fig. 4, SN60 underproduces ^6Li at low metallicity. Our conclusion concerning this ^6Li differs from a previous one [17], namely that the observed ^6Li abundance can be accounted for by HECR spallation. That suggestion was based on the assumption that the composition of the HECR is the same as that of the interstellar medium at low metallicity (i.e. SSz), in which case the high He to CNO ratio for the accelerated particles would indeed yield a high $^6\text{Li}/\text{Be}$. But, as we have seen, such a composition fails energetically to account for the Be abundance and does not yield a constant Be/Fe ratio.

References

- [1] We adopt the convention of using ratios of chemical or isotopic symbols to denote ratios by number of the galactic inventories of the respective elements or isotopes.
- [2] D. K. Duncan et al., ASP Conf. Series 99 (1996) 179.
- [3] J. W. Truran and F. X. Timmes, Nuclei in the Cosmos III (AIP) (1994) 501.
- [4] S. E. Woosley et al., ApJ 356 (1990) 272.
- [5] S. E. Woosley and T. A. Weaver, ApJ (Suppl.) 101 (1995) 181.
- [6] E. Vangioni-Flam et al., ApJ (1996) in press.
- [7] B. D. Fields, K. A. Olive and D. N. Schramm, ApJ 435 (1994) 185.
- [8] R. Ramaty, B. Kozlovsky and R. E. Lingenfelter, ApJ 456 (1996) 525.
- [9] G. Steigman and T. P. Walker, ApJ 385 (1992) L13.
- [10] N. Prantzos, M. Casse and E. Vangioni-Flam, ApJ 403 (1993) 630.

- [11] S. M. Read and V. E. Viola, Atomic Data Nucl. Data Tables 31 (1984) 359.
- [12] W. R. Webber, J. C. Kish and D. A. Schreier, Phys. Rev. C 41 (1990) 547.
- [13] W. R. Webber, private communication 1996.
- [14] J. J. Engelmann et al., A&A 233 (1990) 96.
- [15] N. Lund, Cosmic Abundances of Matter (AIP: New York) (1989) 111.
- [16] S. E. Woosley, N. Langer and T. A. Weaver, ApJ 411 (1993) 823.
- [17] V. V. Smith, D. L. Lambert and P. E. Nissen, ApJ 408 (1993) 262.
- [18] B. Edvardsson et al., A&A 290 (1994) 176.




Figure 1: Number of Be atoms produced per deposited erg. The ambient medium has solar composition scaled with metallicity (SSz). For all curves except SSz the metallicity of the accelerated particles does not change much with $Z/Z_{\odot}$.




Figure 2: B-Be production ratio by accelerated particle interactions. The ambient medium and accelerated particle compositions are as in Fig. 1, except that we do not show the SSz composition. The lowest (dashed) line is for the closed galaxy model.




Figure 3: B isotope production ratio. SN60 achieves values >4 because of its relatively high C/O and sharp cutoff at 50 MeV/n.




Figure 4: ${}^6\text{Li}$ -Be production ratio. SLN represents the observed [17] ${}^6\text{Li}/\text{Li}$ combined with the Li/H and Be/H data.